V.M. Red'kov[*]

Rarita–Schwinger for spin 3/2 field and separation of the variables in static coordinates of de Sitter space, Schrödinger tetrad basis[†]

Institute of Physics, National Academy of Sciences of Belarus

arXiv:1109.4802v1 [math-ph] 22 Sep 2011

Rarita–Schwinger approach to description of a massive spin 3/2 particle is investigated in static coordinates of the de Sitter space-time. The general covariant system, derived from the relevant Lagrangian, is presented as a main wave equation and additional constraints in the form of first order deferential and algebraic relations. With the use of an extended Schrödinger tetrad basis and technique of Wigner D-functions the separation of the variable performed. 16 radial equations reduce to 8 ones through diagonalization of P-inversion operator for spin 3/2 field.

The wave equation in de Sitter space-time for a particle with spin 3/2 can be presented in the form (we adhere notation used in [1]; to the list of references given in [1] one important and much more earliest publication must be added [1]; author apologizes for that inexactness)

$$\Big[\, i\gamma^{\beta}(x) \; (\nabla_{\beta} \; + \; \Gamma_{\beta}(x)) \; - \; m \, \Big] \; \Psi_{\alpha}(x) = 0 \; ,$$

$$\gamma^{\alpha}(x) \; \Psi_{\alpha}(x) = 0 \; , \;\; (\nabla_{\alpha} \; + \; \Gamma_{\alpha}(x)) \; \Psi^{\alpha}(x) = 0 \; . \tag{1}$$

Here the wave function $\Psi_{\alpha}(x)$ is a bispinor with respect to tetrad local rotations and a general covariant vector with respect to general coordinate transformations It will be convenient to use a wave function with tetrad vector index

$$\Psi_{l}(x) = e^{\beta}_{(l)} \; \Psi_{\beta}(x) \; , \quad \Psi_{\beta}(x) = e^{(l)}_{\beta} \; \Psi_{l}(x) \; . \tag{2}$$

Correspondingly the first equation in (1) takes the form

$$\Big[\, i\gamma^{\alpha} \; (\partial_{\alpha} \; + \; \Gamma_{\alpha}(x)) \; \delta^{l}_{k} \; + \; e^{\beta}_{(k)} \; e^{(l)}_{\beta;\alpha}) \; - \; m \, \Big] \; \Psi_{l}(x) = 0 \; ; \tag{3}$$

an addition to connection Γ_{α} in (3) can be presented as vector connection of the type by Tetrode–Weyl–Fock–Ivanenko

$$e^{\beta}_{(k)} \; e^{(l)}_{\beta;\alpha} = (L_{\alpha})_{k}{}^{l} = \frac{1}{2} (J^{ab})_{k}{}^{l} \; e^{\beta}_{(a)} \; (\nabla_{\alpha} e_{(b)\beta}) \; ,$$

---

[*] redkov@dragon.bas-net.by

[†] Chapter 3 in: V.M. Red'kov. Tetrad formalism, spherical symmetry and Schrödinger basis. Publishing House "Belarusian Science", Minsk, 339 pages (2011).

[1] S. Deser, B. Zumino. Consistent supergravity. Phys. Lett. B. 1976. Vol. 62, P. 335 – 337

$$(J^{ab})_k{}^l = \delta^a_k \, g^{bl} \; - \; \delta^b_k \, g^{al} \; . \tag{4}$$

Allowing for (4), eq. (3) is written as

$$[\; i\gamma^{\alpha}(x) \; (\partial_{\alpha} \; + \; B_{\alpha}(x)) \; - \; m \;] \;\; \Psi(x) = 0 \; , \tag{5}$$

$$B_{\alpha}(x) = \Gamma_{\alpha}(x) \otimes I \; + \; I \otimes L_{\alpha}(x) \; . \tag{6}$$

When using the components $\Psi_l(x)$, additional constraints in (1) read

$$\gamma^l \; \Psi_l(x) = 0 \; , \tag{7a}$$

$$\Big[e^{(l)\alpha} \; \partial_{\alpha} \; + \; e^{(l)\alpha}(x) \; + \; e^{(l)\alpha}(x) \; \Gamma_{\alpha}(x) \Big] \; \Psi_l(x) = 0 \; . \tag{7b}$$

We will consider eqs. (6)–(7) in static coordinates

$$dS^2 = (1 - r^2/\rho^2) c^2 \; dt^2 \; - \; \frac{dr^2}{1 - r^2/\rho^2} \; - \; r^2 \; (d\theta^2 \; + \; \sin^2\theta d\phi^2) \; , \tag{8}$$

where ρ stands for the curvature radius. Below we will use dimensionless variables

$$ct/\rho \; \rightarrow \; t, \qquad r/\rho \; \rightarrow \; r \; , \qquad mc\rho/\hbar \; \rightarrow \; m \; . \tag{9}$$

Choosing a diagonal spherical tetrad

$$x^{\alpha} = (t, \; r, \; \theta, \; \phi) \; , \qquad \varphi = 1 - r^2 \; , \qquad \varphi' = d\varphi/dr \; ,$$

$$e^{\alpha}_{(0} = (\varphi^{-1/2}, \; 0, \; 0, \; 0) \; , \qquad e^{\alpha}_{(3)} = (0, \; \varphi^{1/2}, \; 0, \; 0) \; ,$$

$$e^{\alpha}_{(1)} = (0, \; 0, \; 1/r, \; 0) \; , \qquad e^{\alpha}_{(2)} = (0, \; 0, \; 0, \; r^{-1} \sin^{-1}\theta) \; . \tag{10}$$

for $\gamma^{\alpha}(x)$ and $B_{\alpha}(x)$ we get expressions

$$\gamma^{\alpha}(x) = (\; \frac{\gamma^0}{\sqrt{\varphi}} \; , \; \sqrt{\varphi} \, \gamma^3, \; \frac{\gamma^1}{r}, \; \frac{\gamma^2}{r \sin\theta} \;) \; ,$$

$$B_t = \frac{1}{2} \varphi' \; (j^{03} \otimes I + I \otimes J^{03}) \; ,$$

$$\Gamma_r = 0 \; , \;\; \Gamma_{\theta} = \sqrt{\varphi} \; j^{31} \; , \;\; \Gamma_{\phi} = \sqrt{\varphi} \; \sin\theta \; j^{32} \; + \; \cos\theta \; J^{12} \; ,$$

$$L_r = 0 \; , \;\; L_{\theta} = \sqrt{\varphi} \; J^{31} \; , \;\; L_{\phi} = \sqrt{\varphi} \; \sin\theta \; J^{32} \; + \; \cos\theta \; J^{12} \; . \tag{11}$$

Correspondingly, eq. (5) takes the form

$$\Big\{ \; i \frac{\gamma^0}{\sqrt{\varphi}} \partial_0 \; + \; i\sqrt{\varphi} \; [\; \gamma^3 (\partial_r + \frac{1}{r}) \; + \; \frac{\gamma^1 J^{31} \; + \; \gamma^2 J^{32}}{r}$$

$$+ \frac{\varphi'}{2\varphi} \; \gamma^0 \; (j^{03} \otimes I + I \otimes J^{03}) \;] \; + \; \frac{1}{r} \; \Sigma_{\theta,\phi} \; - \; m \; \Big\} \; \Psi(t, r, \theta, \phi) = 0 \; , \tag{12}$$

where $\Sigma_{\theta,\phi}$ is determined by

$$\Sigma_{\theta,\phi} = \; i\gamma^1 \; \partial_{\theta} \; + \; \gamma^2 \; \frac{i\partial_{\phi} \; + (I \otimes iJ^{12} + ij^{12} \otimes I) \cos\theta}{\sin\theta} \; . \tag{13}$$

Further we will use technique extending Schrödinger – Pauli approach to treat problems with spherical symmetry (more details see in [2]). Let us search spherical solutions by diagonalizing operators $\mathbf{J}^2, J_3$ of the total angular momentum. Expressions for these can be found by gauge translation from usual ones in Cartesian tetrad. The latter is known from the theory in flat Minkowski space

$$(\, i\gamma^a \, \partial_a \; - \; m \,) \, \Psi(x) = 0 \; , \; J_i^{Cart} = l_i \; + \; S_i \; ,$$
$$S_1 = iJ^{23} \; , \qquad S_2 = iJ^{31} \; , \qquad S_3 = iJ^{12} \; , \tag{14}$$

where

$$S_i = \frac{1}{2} \, \Sigma_i \otimes I \; + \; I \otimes T_i \; , \qquad T_i = \left| \begin{array}{cc} 0 & 0 \\ 0 & \tau_i \end{array} \right| \; . \tag{15}$$

Wave functions for spin 3/2 particle in two tetrad gages in flat space/ cartesian and spherical, are connected by

$$\Psi^{sph} = S(\theta, \phi) \; \Psi^{Cart} \; , \; \; S = \left| \begin{array}{cc} U_2 & 0 \\ 0 & U_2 \end{array} \right| \otimes \left| \begin{array}{cc} 1 & 0 \\ 0 & O_3 \end{array} \right| \; , \tag{16}$$

where

$$O = \left| \begin{array}{ccc} \cos\theta\cos\phi & \cos\theta\sin\phi & -\sin\theta \\ -\sin\phi & \cos\phi & 0 \\ \sin\theta\cos\phi & \sin\theta\sin\phi & \cos\theta \end{array} \right| , \qquad U_2 = \left| \begin{array}{cc} \cos\theta/2 \; e^{i\phi/2} & \sin\theta/2 \; e^{-i\phi/2} \\ -\sin\theta/2 \; e^{i\phi/2} & \cos\theta/2 \; e^{-i\phi/2} \end{array} \right| .$$

After calculation in accordance with

$$J_i^{Cart} = l_i \; + \; S_i \; , \qquad J_i = S(\theta, \phi) \; J_i^{Cart} \; S^{-1}(\theta, \phi)$$

for components J_i in spherical tetrad basis we get explicit expressions

$$J_1 = l_1 \; + \; S_3 \frac{\sin\phi}{\sin\theta} \; , \qquad J_2 = l_2 \; + \; S_3 \frac{\cos\phi}{\sin\theta} \; , \qquad J_3 = l_3 \; . \tag{17}$$

It is convenient to have matrix $S_3 = iJ^{12}$ as diagonal, but we have

$$S_3 = \frac{1}{2} \left| \begin{array}{cccc} +1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & +1 & 0 \\ 0 & 0 & 0 & -1 \end{array} \right| \otimes I \; + \; I \otimes \left| \begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & +i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right| \quad ,$$

we can reach this by transforming description to the so-called cyclic basis $\tilde{\Psi} = (I \otimes U) \; \Psi$ in vector index

$$\tilde{\Psi} = U \; \Psi \; , \qquad \left| \begin{array}{c} \tilde{\Psi}_{(0)} \\ \tilde{\Psi}_{(1)} \\ \tilde{\Psi}_{(2)} \\ \tilde{\Psi}_{(3)} \end{array} \right| = \left| \begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & -1/\sqrt{2} & i/\sqrt{2} & 0 \\ 0 & 0 & 0 & 1 \\ 0 & +1/\sqrt{2} & i/\sqrt{2} & 0 \end{array} \right| \left| \begin{array}{c} \Psi_{(0)} \\ \Psi_{(1)} \\ \Psi_{(2)} \\ \Psi_{(3)} \end{array} \right| ,$$

$$\Psi = U^{-1}\;\tilde{\Psi}\;, \qquad \left| \begin{array}{c} \Psi_{(0)} \\ \Psi_{(1)} \\ \Psi_{(2)} \\ \Psi_{(3)} \end{array} \right| = \left| \begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & -i/\sqrt{2} & 0 & -i/\sqrt{2} \\ 0 & 0 & 1 & 0 \end{array} \right| \left| \begin{array}{c} \tilde{\Psi}_{(0)} \\ \tilde{\Psi}_{(1)} \\ \tilde{\Psi}_{(2)} \\ \tilde{\Psi}_{(3)} \end{array} \right| ; \qquad (18a)$$

note identities $U^{-1} = U^{+}$, or $(U^{-1})_{kl} = U^{*}_{lk}$.

In the result, eq. (5) assumes the form

$$\Big[\, i\gamma^{\alpha}(x)\;(\partial_{\alpha} \;+\; \Gamma_{\alpha}(x) \otimes I \;+\; I \otimes \tilde{L}_{\alpha}(x)) \;-\; m \,\Big]\;\tilde{\Psi}(x) = 0\;, \qquad (18b)$$

where $\tilde{L}_{\alpha} = U L_{\alpha} U^{-1}$. Additional constraints in cyclic basis will read

$$\gamma^{l}\;\Psi_{l}(x) = \gamma^{l}\;(U^{-1})_{lk}\;\tilde{\Psi}_{k}(x) = U^{*}_{kl}\;\gamma^{l}\;\tilde{\Psi}_{k}(x) = 0\;,$$

$$\Big[\, e^{(l)\alpha}\;\partial_{\alpha} \;+\; e^{(l)\alpha}(x) \;+\; e^{(l)\alpha}(x)\;\Gamma_{\alpha}(x) \,\Big]\;(U^{-1})_{lk}\Psi_{k}(x) = 0\;. \qquad (18c)$$

Transformed generators are

$$\tilde{J}^{ab} = \sigma^{ab} \otimes I \;+\; I \otimes \tilde{j}^{ab}\;, \qquad \tilde{j}^{ab} = U\;j^{ab}\;U^{-1}\;. \qquad (18d)$$

or explicitly

$$i\tilde{J}^{23} = \frac{1}{\sqrt{2}} \left| \begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{array} \right| = \tilde{T}^{1}\;, \qquad i\tilde{J}^{31} = \frac{1}{\sqrt{2}} \left| \begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & +i & 0 & 0-i \\ 0 & 0 & +i & 0 \end{array} \right| = \tilde{T}^{2}\;,$$

$$i\tilde{J}^{12} = \left| \begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & +1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{array} \right| = \tilde{T}^{3}\;, \qquad i\tilde{J}^{03} = \left| \begin{array}{cccc} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right| .$$

Let us turn to eq. (12). Allowing for identities

$$\gamma^{1}\sigma^{31} \;+\; \gamma^{2}\sigma^{32} = \gamma^{3}\;, \qquad \gamma^{0}\sigma^{03} = \frac{1}{2}\,\gamma^{3}$$

and separating a simplifying factor in $\tilde{\Psi}$ according to

$$\tilde{\Psi}(x) = (\epsilon^{-i\epsilon t}/r\varphi^{1/4})\;\tilde{\Phi}(r,\theta,\phi)\;, \qquad (19a)$$

we arrive at the equation for $\tilde{\Phi}(x)$

$$\Big[\, \frac{\gamma^{0}}{\sqrt{\varphi}}\epsilon + i\sqrt{\varphi}(\gamma^{3}\partial_{r} + \frac{\gamma^{1} \otimes \tilde{T}_{2} - \gamma^{2} \otimes \tilde{T}_{1}}{r})$$
$$+\frac{\varphi'}{2\varphi}\gamma^{0}\tilde{J}^{03} + \frac{1}{r}\tilde{\Sigma}_{\theta,\phi} - m \,\Big]\;\tilde{\Phi}(r,\theta,\phi) = 0\;, \qquad (19b)$$

where

$$\tilde{\Sigma}_{\theta,\phi} = i\gamma^{1}\partial_{\theta} \; + \; \gamma^{2}\frac{i\partial_{\phi} \; + \; i\tilde{S}_{3}\cos\theta}{\sin\theta} \; . \tag{19c}$$

Here, the matrix $\tilde{S}_3$ is diagonal: $\tilde{S}_3 = \frac{1}{2}\Sigma_3 \otimes I \; + \; I \otimes \tilde{T}_3$.

There exist 16 possibilities for eigenvector of the matrix S_3; correspondingly we have 16 different eigenvectors for $\vec{J}^2, \; J_3$:

$$D_{-1/2} \left| \begin{array}{c} 1 \\ 0 \\ 0 \\ 0 \end{array} \right| \delta_{l}^{0} \; , \;\; D_{-3/2} \left| \begin{array}{c} 1 \\ 0 \\ 0 \\ 0 \end{array} \right| \delta_{l}^{1} \; , \;\; D_{-1/2} \left| \begin{array}{c} 1 \\ 0 \\ 0 \\ 0 \end{array} \right| \delta_{l}^{2} \; , \;\; D_{+1/2} \left| \begin{array}{c} 1 \\ 0 \\ 0 \\ 0 \end{array} \right| \delta_{l}^{3} \; ,$$

$$D_{+1/2} \left| \begin{array}{c} 0 \\ 1 \\ 0 \\ 0 \end{array} \right| \delta_{l}^{0} \; , \;\; D_{-1/2} \left| \begin{array}{c} 0 \\ 1 \\ 0 \\ 0 \end{array} \right| \delta_{l}^{1} \; , \;\; D_{+1/2} \left| \begin{array}{c} 0 \\ 1 \\ 0 \\ 0 \end{array} \right| \delta_{l}^{2} \; , \;\; D_{+3/2} \left| \begin{array}{c} 0 \\ 1 \\ 0 \\ 0 \end{array} \right| \delta_{l}^{3} \; ;$$

remaining 8 eigenfunctions are constructed in the same manner but with other basis vectors

$$\left| \begin{array}{c} 0 \\ 0 \\ 1 \\ 0 \end{array} \right| \qquad \text{and} \qquad \left| \begin{array}{c} 0 \\ 0 \\ 0 \\ 1 \end{array} \right| \; .$$

Thus, most general substitution for spherical wave solutions with quantum numbers $j, \; m$ can be presented as (bispinor is divided into two -2-spinors)

$$\tilde{\Phi}_{l} \left| \begin{array}{c} \tilde{\xi}_{l} \\ \tilde{\eta}_{l} \end{array} \right| \quad , \;\; \tilde{\xi}_{l} = \left| \begin{array}{c} \tilde{\xi}_{l}^{1} \\ \tilde{\xi}_{l}^{2} \end{array} \right| \quad , \;\; \tilde{\eta}_{l} = \left| \begin{array}{c} \tilde{\eta}_{\dot{1}} \\ \tilde{\eta}_{\dot{2}} \end{array} \right| \quad ,$$

$$\tilde{\xi}_{l} = \left| \begin{array}{c} f_{0} \; \delta_{i}^{0} \; D_{-1/2} \; + \; f_{1} \; \delta_{i}^{1} \; D_{-3/2} \; + \; f_{2} \; \delta_{i}^{2} \; D_{-1/2} \; + \; f_{3} \; \delta_{i}^{3} \; D_{+1/2} \\ g_{0} \; \delta_{i}^{0} \; D_{+1/2} \; + \; g_{1} \; \delta_{i}^{1} \; D_{-1/2} \; + \; g_{2} \; \delta_{i}^{2} \; D_{+1/2} \; + \; g_{3} \; \delta_{i}^{3} \; D_{+3/2} \end{array} \right| \; ; \tag{20a}$$

second 2-spinor $\tilde{\eta}_{i}$ has similar structure but with other radial functions

$$f_{i}(r) \; \Longrightarrow \; h_{i}(r) \; , \;\; g_{i}(r) \; \Longrightarrow \; \nu_{i}(r) \; . \tag{20b}$$

Note that for minimal $j = 1/2$ we must impose restrictions $f_1 = 0, \; h_1 = 0, \; g_3 = 0, \; \nu_3 = 0$.
Taking in mind $\tilde{\Phi}_l = (\tilde{\xi}_i, \; \tilde{\eta}_i)$. eq. (19) is written as

$$\left[\frac{\epsilon}{\sqrt{\varphi}} \; + \; \sqrt{\varphi}(+i\sigma_{3}\partial_{r} \; + \; \frac{\sigma_{1} \otimes \tilde{T}_{2} - \sigma_{2} \otimes \tilde{T}_{1}}{r} + i\frac{\varphi'}{2\varphi} \, \tilde{j}^{03}) \; + \; \frac{1}{r} \, \tilde{\Sigma}_{\theta,\phi} \right] \tilde{\xi} = m \, \tilde{\eta} \; , \tag{21a}$$

$$\left[\frac{\epsilon}{\sqrt{\varphi}} \; + \; \sqrt{\varphi}(-i\sigma_{3}\partial_{r} \; + \; \frac{\sigma_{1} \otimes \tilde{T}_{2} - \sigma_{2} \otimes \tilde{T}_{1}}{r} + i\frac{\varphi'}{2\varphi} \, \tilde{j}^{03}) \; - \; \frac{1}{r} \, \tilde{\Sigma}_{\theta,\phi} \right] \tilde{\eta} = m \, \tilde{\xi} \; , \tag{21b}$$

where

$$\tilde{\Sigma}_{\theta,\phi} = i\sigma_1 \partial_\theta \; + \; \sigma_2 \frac{i\partial_\phi \; + \; (1/2\sigma_3 \otimes I \; + \; I \otimes \tilde{T}_3)\cos\theta}{\sin\theta} \; . \qquad (21c)$$

Additionally to $i\partial_t$, $\vec{J}^2$, J_3 let us diagonalize an operator of spacial inversion. Its form in spherical basis can be found by gauge transformation from the known expression in Cartesian basis

$$\Pi_k^{\;\;l} \; \Psi_l^{Cart}(t, -\vec{r}) \; , \qquad (\Pi_k^{\;\;l}) = \left| \begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{array} \right| \; , \qquad (22a)$$

which results in

$$(\tilde{\Pi}_k^{\;\;l}) = \left| \begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{array} \right| \; . \qquad (22b)$$

An operator of P-inversion in bispinor index will read (in spherical tetrad basis)

$$\Pi = \left| \begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{array} \right| \; . \qquad (23)$$

From eigenvalue equation

$$[\; (\; \Pi \otimes \tilde{\Pi}_k^{\;\;l} \;) \; \hat{P} \;] \; \tilde{\Phi}(r,\theta,\phi) = P \; \tilde{\Phi}(r,\theta,\phi) \qquad (24a)$$

we find two values for parity P and respective linear restrictions on radial functions

$$\nu_0 = \delta \; f_0 \; , \;\; \nu_1 = \delta \; f_3 \; , \;\; \nu_2 = \delta \; f_2 \; , \;\; \nu_3 = \delta \; f_1 \; ,$$

$$h_0 = \delta \; g_0 \; , \;\; h_1 = \delta \; g_3 \; , \;\; h_2 = \delta \; g_2 \; , \;\; h_3 = \delta \; g_1 \; , \qquad (24b)$$

where $\delta = +1$ corresponds to $P = (-1)^{J+1}$ and $\delta = -1$ is referred to $P = (-1)^J$.

Now we are to find radial equations. First let us consider eqs. $(21a, b, c)$ Using relations

$$(\tilde{T}_2)_l^{\;\;k} \; \delta_k^0 = 0 \; , \qquad (\tilde{T}_2)_l^{\;\;k} \; \delta_k^1 = \frac{i}{\sqrt{2}} \; \delta_k^2 \; ,$$

$$(\tilde{T}_2)_l^{\;\;k} \; \delta_k^2 = \frac{i}{\sqrt{2}} \; (\delta_l^3 \; - \; \delta_l^1) \; , \qquad (\tilde{T}_2)_l^{\;\;k} \; \delta_k^3 = -\frac{i}{\sqrt{2}} \; \delta_k^2 \; ,$$

we get

$$(\sigma_1 \otimes \tilde{T}_2)_l^{\;\;k} \; \tilde{\xi}_k = \frac{i}{\sqrt{2}} \left| \begin{array}{c} g_1 \; \delta_l^2 \; D_{-1/2} \; + \; g_2 \; (\delta_l^3 \; - \; \delta_l^1) \; D_{+1/2} \; - \; g_3 \; \delta_l^2 \; D_{+3/2} \\ f_1 \; \delta_l^2 \; D_{-3/2} \; + \; f_2 \; (\delta_l^3 \; - \; \delta_l^1) \; D_{-1/2} - f_3 \; \delta_l^2 \; D_{+1/2} \end{array} \right| \; . \qquad (25)$$

Similarly, allowing for

$$(\tilde{T}_1)_l^{\;\;k} \; \delta_k^0 = 0 \; , \qquad (\tilde{T}_1)_l^{\;\;k} \delta_k^1 = \frac{1}{\sqrt{2}} \delta_k^2 \; ,$$

$$(\tilde{T}_1)_l{}^k \delta^2_k = \frac{i}{\sqrt{2}}(\delta^3_l \ + \ \delta^1_l) \ , \qquad (\tilde{T}_1)_l{}^k \delta^3_k = \frac{1}{\sqrt{2}}\delta^2_k \ ,$$

we obtain

$$-(\sigma_2 \otimes \tilde{T}_1)_l{}^k \ \tilde{\xi}_k = \frac{i}{\sqrt{2}} \left| \begin{array}{c} g_1 \ \delta^2_l \ D_{-1/2} \ + \ g_2 \ (\delta^3_l \ + \ \delta^1_l) \ D_{+1/2} \ + \ g_3 \ \delta^2_l \ D_{+3/2} \\ -f_1 \ \delta^2_l \ D_{-3/2} \ - \ f_2 \ (\delta^3_l \ + \ \delta^1_l) \ D_{-1/2} \ - \ f_3 \ \delta^2_l \ D_{+1/2} \end{array} \right| \ . \tag{26}$$

Therefore, we derive

$$(\sigma_1 \otimes \tilde{T}_2 \ - \ \sigma_2 \otimes \tilde{T}_1)_l{}^k \ \tilde{\xi}_k = i\sqrt{2} \left| \begin{array}{c} +g_1 \ \delta^2_l \ D_{-1/2} \ + \ g_2 \ \delta^3_l \ D_{+1/2} \\ -f_2 \ \delta^1_l \ D_{-1/2} \ - \ g_3 \ \delta^2_l \ D_{+1/2} \end{array} \right| \ . \tag{27}$$

Further, using identities

$$(\tilde{j}^{03})_l{}^k \ \delta^0_k = -\delta^2_l \ , \ \ (\tilde{j}^{03})_l{}^k \ \delta^1_k = 0 \ ,$$

$$(\tilde{j}^{03})_l{}^k \ \delta^2_k = -\delta^0_l \ , \ \ (\tilde{j}^{03})_l{}^k \delta^3_k = 0 \ ,$$

we get

$$i\frac{\varphi'}{2\varphi} \ (\tilde{j}^{03})_l{}^k \ \tilde{\xi}_k = i\frac{\varphi'}{2\varphi} \left| \begin{array}{c} +f_0 \ \delta^2_l \ D_{-1/2} \ + \ f_2 \ \delta^0_l \ D_{-1/2} \\ +g_0 \ \delta^2_l \ D_{+1/2} \ + \ g_2 \ \delta^0_l \ D_{+1/2} \end{array} \right| \ . \tag*{28}$$

To find $\tilde{\Sigma}_{\theta,\phi}$ $(21c)$, one should use the known properties of Wigner function [3]:

$$\partial_\theta \ D_{+1/2} = \frac{1}{2}\left(a \ D_{-1/2} - b \ D_{+3/2}\right) \ ,$$

$$\left[(\sin^{-1}(i\partial_\phi - \frac{1}{2}\cos\theta)\right] \ D_{+1/2} = \frac{1}{2}\left(-a \ D_{-1/2} - b \ D_{+3/2}\right) \ ;$$

$$\partial_\theta \ D_{-1/2} = \frac{1}{2}\left(b \ D_{-3/2} - a \ D_{+1/2}\right) \ ,$$

$$\left[(\sin^{-1}\phi(i\partial + \frac{1}{2}\cos\theta)\right] \ D_{-1/2} = \frac{1}{2}\left(-b \ D_{-3/2} - a \ D_{+1/2}\right) \ ;$$

$$\partial_\theta \ D_{+3/2} = \frac{1}{2}\left(b \ D_{+1/2} - c \ D_{+5/2}\right) \ ,$$

$$\left[(\sin^{-1})(i\partial_\phi - \frac{3}{2}\cos\theta)\right] \ D_{+3/2} = \frac{1}{2}\left(-b \ D_{+1/2} - b \ D_{+5/2}\right) \ ;$$

$$\partial_\theta \ D_{-3/2} = \frac{1}{2}\left(c \ D_{-5/2} - b \ D_{-1/2}\right) \ ,$$

$$\left[(\sin^{-1})(i\partial_\phi + \frac{3}{2}\cos\theta)\right] \ D_{-3/2} = \frac{1}{2}\left(-c \ D_{-5/2} - b \ D_{-1/2}\right) \ , \tag{29a}$$

where

$$a = (j + 1/2) \ , \qquad b = \sqrt{(j - 1/2)(j + 3/2)} \ ,$$

$$c = \sqrt{(j - 3/2)(j + 5/2)} \ . \tag*{29b}$$

So we get

$$(\tilde{\Sigma}_{\theta,\phi}\; \tilde{\xi})_l = i \left| \begin{array}{c} +g_0\; \delta^0_l\; a\; D_{-1/2} \;+\; g_1\; \delta^1_l\; b\; D_{-3/2} \;+\; g_2\; \delta^2_l\; a\; D_{-1/2} \;+\; g_3\; \delta^3_l\; b\; D_{+1/2} \\ -f_0\; \delta^0_l\; a\; D_{+1/2} \;-\; f_1\; \delta^1_l\; b\; D_{-1/2} \;+\; f_2\; \delta^2_l\; a\; D_{+1/2} \;-\; f_3\; \delta^3_l\; b\; D_{+3/2} \end{array} \right| . \qquad (30)$$

For the term $i\sigma_3 \partial_r\; \tilde{\xi}_l$ we obtain

$$i\sigma_3 \partial_r\; \tilde{\xi}_l = i \left| \begin{array}{c} +f'_0\; \delta^0_l\; D_{-1/2} \;+\; f'_1\; \delta^1_l\; D_{-3/2} \;+\; f'_2\; \delta^2_l\; D_{-1/2} \;+\; f'_3\; \delta^3_l\; D_{+1/2} \\ -g'_0\; \delta^0_l\; D_{+1/2} \;-\; g'_1\; \delta^1_l\; D_{-1/2} \;-\; g'_2\; \delta^2_l\; D_{+1/2} \;-\; g'_3\; \delta^3_l\; D_{+3/2} \end{array} \right| . \qquad 31$$

With the use of the above relations we readily derive 8 radial equation

$$\frac{\epsilon}{\sqrt{\varphi}}\; f_0 \;+\; i\sqrt{\varphi}\; \frac{d}{dr}\; f_0 \;-\; i\frac{\varphi'}{2\sqrt{\varphi}}\; f_2 \;+\; i\frac{a}{r}\; g_3 = m\; h_0 \;,$$

$$\frac{\epsilon}{\sqrt{\varphi}}\; g_0 \;-\; i\sqrt{\varphi}\; \frac{d}{dr}\; g_0 \;-\; i\frac{\varphi'}{2\sqrt{\varphi}}\; g_2 \;-\; i\frac{a}{r}\; f_0 = m\; \nu_0 \;. \qquad (32a)$$

$$\frac{\epsilon}{\sqrt{\varphi}}\; f_1 \;+\; i\sqrt{\varphi}\; \frac{d}{dr}\; f_1 \;+\; i\frac{b}{r}\; g_1 = m\; h_1 \;,$$

$$\frac{\epsilon}{\sqrt{\varphi}}\; g_1 \;-\; i\sqrt{\varphi}\; \frac{d}{dr}\; g_1 \;-\; i\frac{\sqrt{2\varphi}}{r}\; f_2 \;-\; i\frac{b}{r}\; f_1 = m\; \nu_1 \;. \qquad (32b)$$

$$\frac{\epsilon}{\sqrt{\varphi}}\; f_2 \;+\; i\sqrt{\varphi}\; \frac{d}{dr}\; f_2 \;-\; i\frac{\varphi'}{2\sqrt{\varphi}}\; f_0 \;+\; i\frac{\sqrt{2\varphi}}{r}\; g_1 \;+\; i\frac{a}{r}\; g_2 = m\; h_2 \;,$$

$$\frac{\epsilon}{\sqrt{\varphi}}\; g_2 \;-\; i\sqrt{\varphi}\; \frac{d}{dr}\; g_2 \;-\; i\frac{\varphi'}{2\sqrt{\varphi}}\; g_0 \;-\; i\frac{\sqrt{2\varphi}}{r}\; f_3 \;-\; i\frac{a}{r}\; f_2 = m\; \nu_2 \;, \qquad (32c)$$

$$\frac{\epsilon}{\sqrt{\varphi}}\; f_3 \;+\; i\sqrt{\varphi}\; \frac{d}{dr}\; f_3 \;+\; i\frac{\sqrt{2\varphi}}{r}\; g_2 \;+\; i\frac{b}{r}\; g_3 = m\; h_3 \;,$$

$$\frac{\epsilon}{\sqrt{\varphi}}\; g_3 \;-\; i\sqrt{\varphi}\; \frac{d}{dr}\; g_3 \;-\; i\frac{b}{r}\; f_3 = m\; \nu_3 \;. \qquad (32d)$$

These 8 equation can be translated to a new (angular) variable ω:

$$r = \sin\omega \;,\; \sqrt{\varphi} = \cos\omega \;,\; \varphi' = -2\sin\omega \;,\;\; \sqrt{\varphi}\frac{d}{dr} = \frac{d}{d\omega} \;.$$

which results in

$$\frac{\epsilon}{\cos\omega}\; f_0 \;+\; i\frac{d}{d\omega}\; f_0 \;+\; i\; \text{tg}\; \omega\; f_2 \;+\; i\frac{a}{\sin\omega}\; g_0 = m\; h_0 \;,$$

$$\frac{\epsilon}{\cos\omega}\; g_0 \;-\; i\frac{d}{d\omega}\; g_0 \;+\; i\; \text{tg}\; \omega\; g_2 \;-\; i\frac{a}{\sin\omega}\; f_0 = m\; \nu_0 \;,$$

$$\frac{\epsilon}{\cos\omega}\; f_1 \;+\; i\frac{d}{d\omega}\; f_1 \;+\; i\frac{b}{\sin\omega}\; g_1 = m\; h_1 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_1 \;-\; i\frac{d}{d\omega}\, g_1 \;-\; i\frac{\sqrt{2}}{\mathrm{tg}\;\omega}\, f_2 \;-\; i\frac{b}{\sin\omega}\, f_1 = m\, \nu_1 \;,$$

$$\frac{\epsilon}{\cos\omega}\, f_2 \;+\; i\frac{d}{d\omega}\, f_2 \;+\; i\;\mathrm{tg}\,\omega\, f_0 \;+\; i\frac{\sqrt{2}}{\mathrm{tg}\;\omega}\, g_1 \;+\; i\frac{a}{\sin\omega}\, g_2 = m\, h_2 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_2 \;-\; i\frac{d}{d\omega}\, g_2 \;+\; i\;\mathrm{tg}\,\omega\, g_0 \;-\; i\frac{\sqrt{2}}{\mathrm{tg}\;\omega}\, f_2 - i\frac{a}{\sin\omega}\, f_2 = m\, \nu_2 \;,$$

$$\frac{\epsilon}{\cos\omega}\, f_3 \;+\; i\frac{d}{d\omega}\, f_3 \;+\; i\frac{\sqrt{2}}{\mathrm{tg}\;\omega}\, g_2 \;+\; i\frac{b}{\sin\omega}\, g_3 = m\, h_3 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_3 \;-\; i\frac{d}{d\omega}\, g_3 \;-\; i\frac{b}{\sin\omega}\, f_3 = m\, \nu_3 \;. \tag{33}$$

Performing formal changes according to

$$f_l(r) \iff h_l(r)\;, \;\; g_l(r) \iff \nu_l(r)$$

and making some small manipulation with signs (compare (21*a*) with (21*b*)), we derive 8 remaining radial equations

$$\frac{\epsilon}{\cos\omega}\, \nu_0 \;+\; i\frac{d}{d\omega}\, \nu_0 \;+\; i\;\mathrm{tg}\,\omega\, \nu_2 \;+\; i\frac{a}{\sin\omega}\, h_0 = m\, g_0 \;,$$

$$\frac{\epsilon}{\cos\omega}\, h_0 \;-\; i\frac{d}{d\omega}\, h_0 \;+\; i\mathrm{tg}\,\omega\, h_2 \;-\; i\frac{a}{\sin\omega}\, \nu_0 = m\, f_0 \;,$$

$$\frac{\epsilon}{\cos\omega}\, \nu_3 \;+\; i\frac{d}{d\omega}\, \nu_3 \;+\; i\frac{b}{\sin\omega}\, h_3 = m\, g_3 \;,$$

$$\frac{\epsilon}{\cos\omega}\, h_3 \;-\; i\frac{d}{d\omega}\, h_3 \;-\; i\frac{\sqrt{2}}{\mathrm{tg}\;\omega}\, \nu_2 \;-\; i\frac{b}{\sin\omega}\, \nu_3 = m\, f_3 \;,$$

$$\frac{\epsilon}{\cos\omega}\, \nu_2 \;+\; i\frac{d}{d\omega}\, \nu_2 \;+\; i\;\mathrm{tg}\,\omega\, \nu_0 \;+\; i\frac{\sqrt{2}}{\mathrm{tg}\;\omega}\, h_3 \;+\; i\frac{a}{\sin\omega}\, h_2 = m\, g_2 \;,$$

$$\frac{\epsilon}{\cos\omega}\, h_2 \;-\; i\frac{d}{d\omega}\, h_2 \;+\; i\;\mathrm{tg}\,\omega\, h_0 \;-\; i\frac{\sqrt{2}}{\mathrm{tg}\;\omega}\, \nu_1 \;-\; i\frac{a}{\sin\omega}\, \nu_2 = m\, f_2 \;,$$

$$\frac{\epsilon}{\cos\omega}\, \nu_1 \;+\; i\frac{d}{d\omega}\, \nu_1 \;+\; i\frac{\sqrt{2}}{\mathrm{tg}\;\omega}\, h_2 \;+\; i\frac{b}{\sin\omega}\, h_1 = m\, g_1 \;,$$

$$\frac{\epsilon}{\cos\omega}\, h_1 \;-\; i\frac{d}{d\omega}\, h_1 \;-\; i\frac{b}{\sin\omega}\, \nu_1 = m\, f_1 \;. \tag{34}$$

This 16-equationm system can be greatly simplified through diagonalization of spacial inversion operator – see linear restrictions (24*b*).

Thus, for parity $(-1)^{J+1}$, we get 8 equations for $f_i(r)$ and $g_i(r)$:

$$\frac{\epsilon}{\cos\omega}\, f_0 \;+\; i\frac{d}{d\omega}\, f_0 \;+\; i\;\mathrm{tg}\,\omega\, f_2 \;+\; i\frac{a}{\sin\omega}\, g_0 = m\, g_0 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_0 \;-\; i\frac{d}{d\omega}\, g_0 \;+\; i \;\text{tg}\,\omega\; g_2 \;-\; i\frac{a}{\sin\omega}\, f_0 = m\, f_0 \;,$$

$$\frac{\epsilon}{\cos\omega}\, f_1 \;+\; i\frac{d}{d\omega}\, f_1 \;+\; i\frac{b}{\sin\omega}\, g_1 = m\, g_3 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_1 \;-\; i\frac{d}{d\omega}\, g_1 \;-\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, f_2 \;-\; i\frac{b}{\sin\omega}\, f_1 = m\, f_3 \;,$$

$$\frac{\epsilon}{\cos\omega}\, f_2 \;+\; i\frac{d}{d\omega}\, f_2 \;+\; i \;\text{tg}\,\omega\; f_0 \;+\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, g_1 \;+\; i\frac{a}{\sin\omega}\, g_2 = m\, g_2 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_2 \;-\; i\frac{d}{d\omega}\, g_2 \;+\; i \;\text{tg}\,\omega\; g_0 \;-\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, f_3 \;-\; i\frac{a}{\sin\omega}\, f_2 = m\, f_2 \;,$$

$$\frac{\epsilon}{\cos\omega}\, f_3 \;+\; i\frac{d}{d\omega}\, f_3 \;+\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, g_2 \;+\; i\frac{b}{\sin\omega}\, g_3 = m\, g_1 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_3 \;-\; i\frac{d}{d\omega}\, g_3 \;-\; i\frac{b}{\sin\omega}\, f_3 = m\, f_1 \;. \qquad (35)$$

For parity (-1)^J, respective equations follow from (35) by changing m into $-m$.

Note that for minimal value $j = 1/2$, the system derived becomes slightly simpler (remember on restrictions $f_1 = 0,\; g_3 = 0,\; b = 0,\; a = 1$):

$$\frac{\epsilon}{\cos\omega}\, f_0 \;+\; i\frac{d}{d\omega}\, f_0 \;+\; i \;\text{tg}\,\omega\; f_2 \;+\; \frac{i}{\sin\omega}\, g_0 = m\, g_0 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_0 \;-\; i\frac{d}{d\omega}\, g_0 \;+\; i \;\text{tg}\,\omega\; g_2 \;-\; \frac{i}{\sin\omega}\, f_0 = m\, f_0 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_1 \;-\; i\frac{d}{d\omega}\, g_1 \;-\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, f_2 = m\, f_3 \;,$$

$$\frac{\epsilon}{\cos\omega}\, f_2 \;+\; i\frac{d}{d\omega}\, f_2 \;+\; i \;\text{tg}\,\omega\; f_0 \;+\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, g_1 \;+\; \frac{i}{\sin\omega}\, g_2 = m\, g_2 \;,$$

$$\frac{\epsilon}{\cos\omega}\, g_2 \;-\; i\frac{d}{d\omega}\, g_2 \;+\; i \;\text{tg}\,\omega\; g_0 \;-\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, f_3 \;-\; \frac{i}{\sin\omega}\, f_2 = m\, f_2 \;,$$

$$\frac{\epsilon}{\cos\omega}\, f_3 \;+\; i\frac{d}{d\omega}\, f_3 \;+\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, g_2 = m\, g_1 \;. \qquad (36)$$

Now we turn to additional constraints ($7a$) and ($7b$)). First consider ($7a$); it can be written as (expression for U see in ($18b$))

$$\tilde{\xi}_0(x) \;+\; \sigma_i\, U^{-1}_{ij}\, \tilde{\xi}_j(x) = 0 \;, \qquad \tilde{\eta}_0(x) \;-\; \sigma_i\, U^{-1}_{ij}\, \tilde{\eta}_j(x) = 0 \;. \qquad 37a$$

From the first equation in ($37a$) it follows

$$f_0 D_{-1/2} \; + \; \frac{1}{\sqrt{2}} \, (-g_1 \; D_{-1/2} + g_3 \; D_{+3/2}) \; +$$

$$f_2 \; D_{-1/2} \; - \; \frac{1}{\sqrt{2}} \, (+g_1 \; D_{-1/2} + g_3 \; D_{+3/2}) = 0 \; ,$$

$$g_0 \; D_{+1/2} \; + \; \frac{1}{\sqrt{2}} \, (-f_1 \; D_{-3/2} + f_3 \; D_{+1/2}) \; -$$

$$g_2 \; D_{+1/2} \; + \; \frac{1}{\sqrt{2}} (+f_1 \; D_{-3/2} + f_3 \; D_{+1/2}) = 0 \; , \tag{37b}$$

which results in two algebraic constraints

$$f_0 \; - \; \sqrt{2} \, g_1 \; + \; f_2 = 0 \; , \qquad g_0 \; + \; \sqrt{2} \, f_3 \; - \; g_2 = 0 \; . \tag{38a}$$

In the same manner, from second equation in $(37a)$ it follows

$$h_0 \; + \; \sqrt{2} \, \nu_1 \; - \; h_2 = 0 \; , \qquad \nu_0 \; - \; \sqrt{2} \, h_3 \; + \; \nu_2 = 0 \; . \tag{38b}$$

Allowing for $(24b)$, we see that at both values of parity, (37) and (38) give the same two relationships

$$f_0 \; - \; \sqrt{2} \, g_1 \; + \; f_2 = 0 \; , \qquad g_0 \; + \; \sqrt{2} \, f_3 \; - \; g_2 = 0 \; . \tag{39}$$

Note that these two relations are valid at minimal $j = 1/2$ as well.

Now consider the second additional condition $(7b)$; it can be presented as

$$(\nabla_\alpha + \Gamma_\alpha) \; (e^{(l)\alpha} \Psi_{(l)}) = [\; e^{(l)\alpha} \; \partial_\alpha \; + \; e^{(l)\alpha}_{;\alpha}(x) \; + \; e^{(l)\alpha}(x) \; \Gamma_\alpha(x)] \; \Psi_l(x) = 0 \; , \tag{40}$$

where $\Psi_l(x) = U^{-1}_{lk} \; \tilde{\Psi}_k$. Allowing for

$$U^{-1}_{lk} \delta^0_k = \delta^0_l \; , \; U^{-1}_{lk} \delta^1_k = \frac{1}{\sqrt{2}} \, (-\delta^1_l \; - \; i\delta^2_l) \; ,$$

$$U^{-1}_{lk} \delta^2_k = \delta^3_l \; , \;\; U^{-1}_{lk} \delta^3_k = \frac{1}{\sqrt{2}} \, (+\delta^1_l \; - \; i\delta^2_l) \; ,$$

we find 2-spinors $\xi_l(x) = U^{-1}_{lk} \; \tilde{\xi}_k$ and $\eta_l(x) = U^{-1}_{lk} \; \tilde{\eta}_k$:

$$(\xi_l) =$$

$$\left| \begin{array}{c} f_0 \; \delta^0_l \; D_{-1/2} \; + \; f_1 \frac{1}{\sqrt{2}} \, (-\delta^1_l - i\delta^2_l) \; D_{-3/2} \; + \; f_2 \; \delta^3_l \; D_{-1/2} \; + \; f_3 \frac{1}{\sqrt{2}} \, (+\delta^1_l \; - \; i\delta^2_l) \; D_{+1/2} \\ g_0 \; \delta^0_l \; D_{+1/2} \; + \; g_1 \frac{1}{\sqrt{2}} \, (-\delta^1_l \; - \; i\delta^2_l) \; D_{-1/2} \; + \; g_2 \; \delta^3_l \; D_{+1/2} \; + \; g_3 \frac{1}{\sqrt{2}} \, (+\delta^1_l \; - \; i\delta^2_l) \; D_{+3/2} \end{array} \right| \; . \tag{41}$$

Expression for (η_l) will be similar but with other radial functions $h_i(r)$ and $\nu_i(r)$. Further, we find

$$e^{(l)\alpha} \partial_\alpha \; \Psi_l = e^{(0)0} \; \partial_0 \Psi_0 + e^{(3)r} \; \partial_r \Psi_3 + e^{(1)\theta} \; \partial_\theta \Psi_1 + e^{(2)\phi} \; \partial_\phi \Psi_2$$

$$= \frac{-i\epsilon}{\sqrt{\varphi}} \left| \begin{array}{l} f_0\ D_{-1/2} \\ g_0\ D_{+1/2} \\ h_0\ D_{-1/2} \\ \nu_0\ D_{+1/2} \end{array} \right| - \frac{1}{\sqrt{2}r}\partial_\theta \left| \begin{array}{l} -f_1\ D_{-3/2} + f_3\ D_{+1/2} \\ -g_1\ D_{-1/2} + g_3\ D_{+3/2} \\ -h_1\ D_{-3/2} + h_3\ D_{+1/2} \\ -\nu_1\ D_{-1/2} + \nu_3\ D_{+3/2} \end{array} \right|$$

$$-\sqrt{\varphi}\partial_r \left| \begin{array}{l} f_2\ D_{-1/2} \\ g_2\ D_{+1/2} \\ h_2\ D_{-1/2} \\ \nu_2\ D_{+1/2} \end{array} \right| + \frac{i\partial_\phi}{\sqrt{2}r\sin\theta} \left| \begin{array}{l} +f_1\ D_{-3/2} + f_3\ D_{+1/2} \\ +g_1\ D_{-1/2} + g_3\ D_{+3/2} \\ +h_1\ D_{-3/2} + h_3\ D_{+1/2} \\ +\nu_1\ D_{-1/2} + \nu_3\ D_{+3/2} \end{array} \right| . \tag{42}$$

With the help of identities

$$e^{(0)\alpha}_{\ ;\alpha} = 0 \ , \quad e^{(1)\alpha}_{\ ;\alpha} = -\frac{1}{r}\ \mathrm{ctg}\ \theta \ , \quad e^{(2)\alpha}_{\ ;\alpha} = 0 \ , \quad e^{(3)\alpha}_{\ ;\alpha} = -\sqrt{\varphi}(\frac{2}{r} + \frac{\varphi'}{2\varphi}) \ ,$$

we get

$$e^{(l)\alpha}_{\ ;\alpha}\ \Psi_l = \frac{\mathrm{ctg}\ \theta}{\sqrt{2}r} \left| \begin{array}{l} f_1\ D_{-3/2} - f_3\ D_{+1/2} \\ g_1\ D_{-1/2} - g_3\ D_{+3/2} \\ h_1\ D_{-3/2} - h_3\ D_{+1/2} \\ \nu_1\ D_{-1/2} - \nu_3\ D_{+3/2} \end{array} \right| - \sqrt{\varphi}(\frac{2}{r} + \frac{\varphi'}{2\varphi}) \left| \begin{array}{l} f_2\ D_{-1/2} \\ g_2\ D_{+1/2} \\ h_2\ D_{-1/2} \\ \nu_2\ D_{+1/2} \end{array} \right| . \tag{43}$$

As for the term $e^{(l)\alpha}\ \Gamma_\alpha\ \Psi_l$; taking into account

$$e^{(l)\alpha}\ \Gamma_\alpha\ \Psi_l = e^{(0)0}\ \Gamma_0\ \Psi_0 + e^{(3)r}\ \Gamma_r\ \Psi_3 + e^{(1)\theta}\ \Gamma_\theta\ \Psi_1 + e^{(2)\phi}\ \Gamma_\phi\ \Psi_2 =$$

$$\frac{\varphi'}{2\sqrt{\varphi}}\ \sigma^{03}\Psi_0 \ - \ \frac{\sqrt{\varphi}}{r}\ (\sigma^{31}\Psi_1 \ + \ \sigma^{32}\Psi_2) - \frac{\mathrm{ctg}\ \theta}{r}\ \sigma^{21}\Psi_2 \tag{44a}$$

after simple calculation we get

$$e^{(l)\alpha}\Gamma_\alpha\ \Psi_l$$

$$= \frac{\varphi'}{4\sqrt{\varphi}} \left| \begin{array}{r} f_0\ D_{-1/2} \\ -g_0\ D_{+1/2} \\ -h_0\ D_{-1/2} \\ \nu_0\ D_{+1/2} \end{array} \right| - \frac{\sqrt{\varphi}}{\sqrt{2}r} \left| \begin{array}{l} g_1\ D_{-1/2} \\ f_3\ D_{+1/2} \\ \nu_1\ D_{-1/2} \\ h_3\ D_{+1/2} \end{array} \right| + \frac{\mathrm{ctg}\ \theta}{\sqrt{2}r} \left| \begin{array}{l} -f_1\ D_{-3/2} - f_3\ D_{+1/2} \\ +g_1\ D_{-1/2} + g_3\ D_{+3/2} \\ -h_1\ D_{-3/2} - h_3\ D_{+1/2} \\ +\nu_1\ D_{-1/2} + \nu_3\ D_{+3/2} \end{array} \right| . \tag{44b}$$

Summing (42), (43), and (44b), we obtain for (40)

$$\frac{-i\epsilon}{\sqrt{\varphi}} \left| \begin{array}{l} f_0\ D_{-1/2} \\ g_0\ D_{+1/2} \\ h_0\ D_{-1/2} \\ \nu_0\ D_{+1/2} \end{array} \right| + \frac{\varphi'}{4\sqrt{\varphi}} \left| \begin{array}{r} f_0\ D_{-1/2} \\ -g_0\ D_{+1/2} \\ -h_0\ D_{-1/2} \\ \nu_0\ D_{+1/2} \end{array} \right|$$

$$-\sqrt{\varphi}(\partial_r + \frac{2}{r} + \frac{\varphi'}{2\varphi}) \left| \begin{array}{l} f_2\ D_{-1/2} \\ g_2\ D_{+1/2} \\ h_2\ D_{-1/2} \\ \nu_2\ D_{+1/2} \end{array} \right| - \frac{\sqrt{\varphi}}{\sqrt{2}r} \left| \begin{array}{l} g_1\ D_{-1/2} \\ f_3\ D_{+1/2} \\ \nu_1\ D_{-1/2} \\ h_3\ D_{+1/2} \end{array} \right|$$

$$
-\frac{1}{\sqrt{2}\;r}\partial_\theta \left| \begin{array}{c} -f_1\;D_{-3/2}+f_3\;D_{+1/2} \\ -g_1\;D_{-1/2}+g_3\;D_{+3/2} \\ -h_1\;D_{-3/2}+h_3\;D_{+1/2} \\ -\nu_1\;D_{-1/2}+\nu_3\;D_{+3/2} \end{array} \right|
$$

$$
-\frac{1}{\sqrt{2}r} \left| \begin{array}{ccc} f_1\;\frac{i\partial_\phi+3/2\cos\theta}{\sin\theta}D_{-3/2} & + & f_3\;\frac{i\partial_\phi-1/2\cos\theta}{\sin\theta}D_{+1/2} \\ g_1\;\frac{i\partial_\phi+1/2\cos\theta}{\sin\theta}D_{-1/2} & + & g_3\;\frac{i\partial_\phi-3/2\cos\theta}{\sin\theta}D_{+3/2} \\ h_1\;\frac{i\partial_\phi+3/2\cos\theta}{\sin\theta}D_{-3/2} & + & h_3\;\frac{i\partial_\phi-1/2\cos\theta}{\sin\theta}D_{+3/2} \\ \nu_1\;\frac{i\partial_\phi+1/2\cos\theta}{\sin\theta}D_{-1/2} & + & \nu_3\;\frac{i\partial_\phi-3/2\cos\theta}{\sin\theta}D_{+3/2} \end{array} \right| = 0\;.
$$

From whence, transforming two last terms with the help of (29), we get

$$
D_\beta\;\Psi^\beta = \frac{-i\epsilon}{\sqrt{\varphi}} \left| \begin{array}{c} f_0\;D_{-1/2} \\ g_0\;D_{+1/2} \\ h_0\;D_{-1/2} \\ \nu_0\;D_{+1/2} \end{array} \right| + \frac{\varphi'}{4\sqrt{\varphi}} \left| \begin{array}{r} f_0\;D_{-1/2} \\ -g_0\;D_{+1/2} \\ -h_0\;D_{-1/2} \\ \nu_0\;D_{+1/2} \end{array} \right|
$$

$$
-\sqrt{\varphi}(\partial_r + \frac{2}{r} + \frac{\varphi'}{2\varphi}) \left| \begin{array}{c} f_2\;D_{-1/2} \\ g_2\;D_{+1/2} \\ h_2\;D_{-1/2} \\ \nu_2\;D_{+1/2} \end{array} \right|
$$

$$
-\frac{\sqrt{\varphi}}{\sqrt{2}r} \left| \begin{array}{c} g_1\;D_{-1/2} \\ f_3\;D_{+1/2} \\ \nu_1\;D_{-1/2} \\ h_3\;D_{+1/2} \end{array} \right| - \frac{1}{\sqrt{2}\;r} \left| \begin{array}{c} (b\;f_1+a\;f_3)\;D_{-1/2} \\ (a\;g_1+b\;g_3)\;D_{-1/2} \\ (b\;h_1+a\;h_3)\;D_{-1/2} \\ (a\;\nu_1+b\;\mu_3)\;D_{-1/2} \end{array} \right| . \tag{45}
$$

Thus, we derive four differential relations – transform them to the variable ω and take into account $(19a)$):

$$
\frac{-i\epsilon}{\cos\omega}\;f_0 \;-\; \frac{\mathrm{tg}\;\omega}{2}\;f_0 \;-\; \frac{d}{d\omega}\;f_2 \;-\; \frac{1}{\sqrt{2}\;\mathrm{tg}\;\omega}\;g_1 \;-\; \frac{1}{\sqrt{2}\sin\omega}\;(b\;f_1+a\;f_3)=0\;,
$$

$$
\frac{-i\epsilon}{\cos\omega}\;g_0 \;+\; \frac{\mathrm{tg}\;\omega}{2}\;g_0 \;-\; \frac{d}{d\omega}\;g_2 \;-\; \frac{1}{\sqrt{2}\;\mathrm{tg}\;\omega}\;f_3 \;-\; \frac{1}{\sqrt{2}\sin\omega}\;(a\;g_1+b\;g_3)=0\;,
$$

$$
\frac{-i\epsilon}{\cos\omega}\;h_0 \;+\; \frac{\mathrm{tg}\;\omega}{2}\;h_0 \;-\; \frac{d}{d\omega}\;h_2 \;-\; \frac{1}{\sqrt{2}\;\mathrm{tg}\;\omega}\;\nu_1 \;-\; \frac{1}{\sqrt{2}\sin\omega}\;(b\;h_1+a\;h_3)=0\;,
$$

$$
\frac{-i\epsilon}{\cos\omega}\;\nu_0 \;-\; \frac{\mathrm{tg}\;\omega}{2}\;\nu_0 \;-\; \frac{d}{d\omega}\;\nu_2 \;-\; \frac{1}{\sqrt{2}\;\mathrm{tg}\;\omega}\;h_3 \;-\; \frac{1}{\sqrt{2}\sin\omega}\;(a\;\nu_1+b\;\nu_3)=0\;. \tag{46a}
$$

Note that two first equations in $(46a)$ include only f_l and g_l, whereas two las t contain h_l and ν_l. When taking into account restrictions (24), from $(46a)$ it remain only two first relations

$$
\frac{-i\epsilon}{\cos\theta}\;f_0 \;-\; \frac{\mathrm{tg}\;\omega}{2}\;f_0 \;-\; \frac{d}{d\omega}\;f_2 \;-\; \frac{1}{\sqrt{2}\;\mathrm{tg}\;\omega}\;g_1 \;-\; \frac{1}{\sqrt{2}\sin\omega}\;(b\;f_1+a\;f_3)=0\;,
$$

$$\frac{-i\epsilon}{\cos\theta}\, g_0 \;+\; \frac{\text{tg}\,\omega}{2}\, g_0 \;-\; \frac{d}{d\omega}\, g_2 \;-\; \frac{1}{\sqrt{2}\,\text{tg}\,\omega}\, f_3 \;-\; \frac{1}{\sqrt{2}\sin\omega}\,(a\, g_1 + b\, g_3) = 0\;. \tag{46b}$$

For minimal $j = 1/2$ these equation become slightly simpler

$$\frac{-i\epsilon}{\cos\theta}\, f_0 \;-\; \frac{\text{tg}\,\omega}{2}\, f_0 \;-\; \frac{d}{d\omega}\, f_2 \;-\; \frac{1}{\sqrt{2}\,\text{tg}\,\omega}\, g_1 \;-\; \frac{1}{\sqrt{2}\sin\omega}\, f_3 = 0\;,$$

$$\frac{-i\epsilon}{\cos\theta}\, g_0 \;+\; \frac{\text{tg}\,\omega}{2}\, g_0 \;-\; \frac{d}{d\omega}\, g_2 \;-\; \frac{1}{\sqrt{2}\,\text{tg}\,\omega}\, f_3 \;-\; \frac{1}{\sqrt{2}\sin\omega}\, g_1 = 0\;. \tag{47}$$

Let us collect results together

after separation the variables in the equations describing the massive particle with spin 3/2 in de Sitter space-time, the problem is reduced to

1) 8 main radial equations at the parity $(-1)^{J+1}$ (for other parity $(-1)^J$, parameter m should be changed into $-m$)

$$\frac{\epsilon}{\cos\omega}\, f_0 \;+\; i\frac{d}{d\omega}\, f_0 \;+\; i\;\text{tg}\,\omega\; f_2 \;+\; i\frac{a}{\sin\omega}\, g_0 = m\; g_0\;,$$

$$\frac{\epsilon}{\cos\omega}\, g_0 \;-\; i\frac{d}{d\omega}\, g_0 \;+\; i\;\text{tg}\,\omega\; g_2 \;-\; i\frac{a}{\sin\omega}\, f_0 = m\; f_0\;,$$

$$\frac{\epsilon}{\cos\omega}\, f_1 \;+\; i\frac{d}{d\omega}\, f_1 \;+\; i\frac{b}{\sin\omega}\, g_1 = m\; g_3\;,$$

$$\frac{\epsilon}{\cos\omega}\, g_1 \;-\; i\frac{d}{d\omega}\, g_1 \;-\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, f_2 \;-\; i\frac{b}{\sin\omega}\, f_1 = m\; f_3\;,$$

$$\frac{\epsilon}{\cos\omega}\, f_2 \;+\; i\frac{d}{d\omega}\, f_2 \;+\; i\;\text{tg}\,\omega\; f_0 \;+\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, g_1 \;+\; i\frac{a}{\sin\omega}\, g_2 = m\; g_2\;,$$

$$\frac{\epsilon}{\cos\omega}\, g_2 \;-\; i\frac{d}{d\omega}\, g_2 \;+\; i\;\text{tg}\,\omega\; g_0 \;-\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, f_3 \;-\; i\frac{a}{\sin\omega}\, f_2 = m\; f_2\;,$$

$$\frac{\epsilon}{\cos\omega}\, f_3 \;+\; i\frac{d}{d\omega}\, f_3 \;+\; i\frac{\sqrt{2}}{\text{tg}\,\omega}\, g_2 \;+\; i\frac{b}{\sin\omega}\, g_3 = m\; g_1\;,$$

$$\frac{\epsilon}{\cos\omega}\, g_3 \;-\; i\frac{d}{d\omega}\, g_3 \;-\; i\frac{b}{\sin\omega}\, f_3 = m\; f_1\;; \tag{48a}$$

2) 2 algebraic constraints (the same for both parities)

$$g_1 = \frac{f_2 + f_0}{\sqrt{2}}\;, \qquad f_3 = \frac{g_2 - g_0}{\sqrt{2}}\;; \tag{48b}$$

and two differential constraints (the same for both parities)

$$\frac{-i\epsilon}{\cos\theta}\, f_0 \;-\; \frac{\text{tg}\,\omega}{2}\, f_0 \;-\; \frac{d}{d\omega}\, f_2 \;-\; \frac{1}{\sqrt{2}\,\text{tg}\,\omega}\, g_1 \;-\; \frac{1}{\sqrt{2}\sin\omega}\,(b\, f_1 + a\, f_3) = 0\;,$$

$$\frac{-i\epsilon}{\cos\theta}\, g_0 \;+\; \frac{\mathrm{tg}\,\omega}{2}\, g_0 \;-\; \frac{d}{d\omega}\, g_2 \;-\; \frac{1}{\sqrt{2}\;\mathrm{tg}\,\omega}\, f_3 \;-\; \frac{1}{\sqrt{2}\sin\omega}\,(a\, g_1 + b\, g_3) = 0\ . \tag{48c}$$

Note that the same technique can be used to separate the variables in any space-time with spherical symmetry, also in presence of any external electromagnetic field with spherical symmetry (in this case additional constrains should be extended – see in [1]).

References

[1] V.M. Red'kov. Particle with spin S = 3/2 in Riemannian space-time. 12 pages. arXiv:1109.3871v1 [math-ph] 18 Sep 2011

[2] V.M. Red'kov. Tetrad formalism, spherical symmetry and Schrödinger basis. Publishing House "Belarusian Science", Minsk, 339 pages, 2011 (in Russian)

[3] D.A. Varshalovich, A.N. Moskalev, V.K. Hersonskiy. Quantum theory of angular moment. Nauka, Leningrad, 1975 (in Russian)